

# ON PROJECTIVE VARIETIES WITH NEF ANTICANONICAL DIVISORS

QI ZHANG

## §1. Introduction

The aim of this note is to prove a structure theorem for projective varieties with nef anticanonical divisors (the Main Theorem). In [18], we showed that if  $X$  is smooth and  $-K_X$  is nef, then the Albanese map  $\text{Alb}_X : X \rightarrow \text{Alb}(X)$  is surjective and has connected fibers (i.e., it is a fiberspace map). In this note we apply the techniques which have been developed in [2],[14] and [19] to prove the following;

**Main Theorem.** *Let  $X$  be a projective variety and  $D$  an effective  $\mathbb{Q}$ -divisor on  $X$  such that the pair  $(X, D)$  is log canonical [10] and  $-(K_X + D)$  is nef. Let  $f : X \dashrightarrow Y$  be a dominant rational map, where  $Y$  is a smooth variety. Then either*

- (1)  $Y$  is uniruled; or
- (2) *The Kodaira dimension  $\kappa(Y) = 0$  Moreover in this case,  $f$  is semistable in codimension 1 (see Definition 1).*

*Remark 1:* In fact, we only need to assume that  $(X, D)$  is log canonical along the general fibers of  $f$ . Also, in case (2), we actually have  $\kappa_\sigma(Y) = 0$  [14]. Moreover, our proof also shows that in this case, every component of  $D$  dominates  $Y$ . This fact was proved by McKernan-Prokhorov in [20, Lemma 10.2] under the assumption of the Log Minimal Model Conjecture.

*Remark 2:* When  $X$  is smooth and  $f$  is a morphism, the Main Theorem was essentially proved in [18] by using the relative deformation theory and the mod  $p$  reduction methods. However, the proof we present here is new and is completely different in nature.

Applying the Main Theorem to the maximal rationally connected fibration of  $X$  [3],[11] and using a result of Graber-Harris-Starr [7], we obtain the following:

**Corollary 1.** *Let  $X$  be a projective variety and  $D$  an effective  $\mathbb{Q}$ -divisor such that the pair  $(X, D)$  is log canonical and  $-(K_X + D)$  is nef. Then there exists a dominant rational map  $f : X \dashrightarrow Y$  such that*

- (1)  *$f$  is a fibration map which is also semistable in codimension one.*
- (2) *The general fibers of  $f$  are rationally connected.*
- (3)  *$Y$  is smooth with  $\kappa(Y) = 0$ .*

*Remark 3:* In the statement of Corollary 1, I believe that  $Y$  should be birational to a variety  $W$  with log canonical singularities and  $K_W$  being numerically trivial.

**Corollary 2.** *Let  $X$  be a projective variety and  $D$  an effective  $\mathbb{Q}$ -divisor such that the pair  $(X, D)$  is log canonical and  $-(K_X + D)$  is nef. Then the Albanese map (from any smooth model of  $X$ )  $\text{Alb}_X : X \dashrightarrow \text{Alb}(X)$  is a fibration map which is also semistable in codimension one.*

*Remark 4:* A conjecture of Demailly-Peternell-Schneider [6] predicts that when  $X$  is smooth and  $-K_X$  is nef, the Albanese map should be a smooth morphism. When  $\dim X = 3$ , this has been verified by Peternell-Serrano [15]. Also, Campana proved that when  $X$  is a special variety, its Albanese map is multiple fibers free in codimension one [4].

**Corollary 3.** *Let  $X$  be a projective variety and  $D$  an effective  $\mathbb{Q}$ -divisor such that the pair  $(X, D)$  is log canonical and  $-(K_X + D)$  is nef. Let  $f : X \dashrightarrow Y$  be the nef reduction of  $-(K_X + D)$  [5],[16]. Then  $Y$  is uniruled.*

*Remark 5:* The notion of special varieties was introduced and studied by F. Campana in [4]. He also conjectured that compact Kähler manifolds with  $-K_X$  nef are special. S. Lu [12] proved the conjecture for projective varieties. In particular, if  $X$  is a projective variety with  $-K_X$  nef and if there is a surjective map  $X \rightarrow Y$ . Then  $\kappa(Y) \leq 0$ . Our focus however, is on the uniruledness of  $Y$ .

*Remark 6:* When  $\dim X \leq 4$  and  $-K_X$  is nef, we can easily derive some (rough) classification results for those varieties from Corollaries 1-3. A complete classification of such threefolds was given in [21].

The tools we shall use in the proof of the Main Theorem are the theory of weak (semi) positivity of the direct images of (log) relative dualizing sheaves  $f_*(K_{X/Y} + \Delta)$  (which has been developed by Fujita, Kawamata, Kollár, Viehweg and others) plus the notion of the divisorial Zariski decomposition [1],[14]. Also, some results in [2] are crucial to the proof.

## §2. Proofs

Before we start to prove the Main Theorem and Corollaries 1-3, let us first give the related definition.

**Definition 1.** *Let  $f : X \dashrightarrow Y$  be a dominant rational map, where  $X$  and  $Y$  are normal proper varieties. We say that  $f$  is semistable in codimension one if it satisfies the following condition :*

*Let  $\pi : Z \rightarrow X$  be a birational morphism which is an eliminating the indeterminacy of  $f$ . Let  $P$  be a prime divisor on  $Y$ . If  $D$  is a non-reduced component of  $g^*(P)$ , where  $g = f \circ \pi$ . Then either  $\text{codim}(g(D)) \geq 2$  or  $D$  is  $\pi$ -exceptional.*

We work over the complex number field  $\mathbb{C}$  in this paper.

*Proof of the Main Theorem.* The proof is quite similar to the proof of Proposition 1 in [19].

Eliminating the indeterminacy of  $f$  and taking a log resolution, we have a smooth projective variety  $Z$ , and the surjective morphisms  $g : Z \rightarrow Y$  and  $\pi : Z \rightarrow X$ ,

$$\begin{array}{ccc} Z & \xrightarrow{\pi} & X \\ \downarrow g & & \\ Y & & \end{array}$$

such that  $K_Z = \pi^*(K_X + D) + \sum e_i E_i$  with  $e_i \geq -1$ , where  $\sum E_i$  is a divisor with normal crossing.

Let  $\delta L$  be an ample  $\mathbb{Q}$ -divisor (with small  $\delta > 0$ ) on  $Z$ . Since  $-(K_X + D)$  is nef,  $-\pi^*(K_X + D) + \delta L$  is ample. Let  $H$  be an ample  $\mathbb{Q}$ -divisor on  $Y$  such that  $A = -\pi^*(K_X + D) + \delta L - H$  is again an ample  $\mathbb{Q}$ -effective divisor. We may also assume that  $\text{Supp}(A + \sum E_i)$  is a divisor with simple normal crossing. Let  $\Delta = A + \sum \epsilon_i E_i$ , where  $\epsilon_i E_i = -e_i E_i$  if  $e_i < 0$  and  $\epsilon_i E_i = \{-e_i\} E_i$  if  $e_i \geq 0$ . We may assume that the pair  $(X, \Delta)$  is log canonical. We have

$$K_{Z/Y} + \Delta \sim_{\mathbb{Q}} \sum m_i E_i - g^*(K_Y + H) + \delta L$$

where  $m_i = \lceil e_i \rceil$  (for  $e_i \geq 0$ ) are non-negative integers (where  $\{, \}$  is the fractional part and  $\lceil, \rceil$  is the round up).

By the same arguments as in [19] (using the semistable reduction theorem, the covering trick and the flattening of  $g$ ), there exist a finite morphism  $p : Y' \rightarrow Y$  and the induced morphism  $g' : Z' \rightarrow Y'$  from a desingularization

$Z' \rightarrow Z \times_Y Y'$  such that it is semistable over  $Y' \setminus B$  with  $\text{codim}(B) \geq 2$ . Let  $Z' \rightarrow Z$  be the induced morphism.

$$\begin{array}{ccccc} Z' & \xrightarrow{q} & Z & \xrightarrow{\pi} & X \\ g' \downarrow & & g \downarrow & & \\ Y' & \xrightarrow{p} & Y & & \end{array}$$

By the ramification formula and the fact that the relative dualizing sheaves are stable under the base change. By [19] we have

$$K_{Z'/Y'} + \Delta' \sim_Q \sum n_{k'} E'_{k'} - g'^* p^*(K_Y + H) + q^*(\delta L) - V + G$$

where

- (1)  $\sum n_{k'} E'_{k'}$  is Cartier and  $\text{Supp}(\sum E'_{k'})$  is  $q \circ \pi$ -exceptional.
- (2)  $V$  is an effective Cartier divisor which is  $g'$ -vertical ( $V$  contains the divisors from the non-reduced components of the fibers of  $g$ ).
- (3)  $G$  is also Cartier and  $q \circ \pi$ -exceptional ( $\text{codim}_{g'}(G) \geq 2$ ).
- (4) If  $E'_{k'}$  is  $g'$ -horizontal, then  $n_{k'} \geq 0$  ( $g'$ -horizontal divisors are not the ramification divisors of  $q$ ).
- (5)  $(Z', \Delta')$  is log canonical.
- (6)  $p^*H = 2H'$  where  $H'$  is an ample Cartier divisor.
- (7) If a divisor  $D$  on  $Z'$  with  $\text{codim}(g'(D)) \geq 2$ , then  $D$  is  $q \circ \pi$ -exceptional.

The following weak positivity theorem is a generalization of the results due to Kawamata [8], [9] and Viehweg [17], and it was proved first by F. Campana [4, Theorem 4.13] (see also [12]).

**Lemma 1.** *Let  $U$  be a smooth projective variety and  $M$  be an effective divisor such that the pair  $(U, M)$  is log canonical. Let  $\phi : U \rightarrow W$  be a surjective morphism, where  $W$  is smooth. Then  $\phi_*(m(K_{U/W} + D))$  is torsion free and weakly positive for any  $m \in \mathbb{Z}_{>0}$  such that  $m(K_{U/W} + D)$  is Cartier.*

Let us choose a positive integer  $m > 0$  such that  $m(\delta L)$  is Cartier.

Since all the  $g'$ -horizontal divisors  $E'_{k'}$  and  $q^*(m\delta L)$  have non negative coefficients,

$$g'_*(m(\sum n_{k'} E''_{k'} - g'^* p^*(K_Y) - g'^*(2H') + q^*(\delta L) - V + G))$$

is a non-zero sheaf. Let

$$\omega = m(\sum n_{k'} E'_{k'} - g'^* p^*(K_Y) + q^*(\delta L) - V + G)$$

By Lemma 1,  $g'_*(\omega) \otimes \mathcal{O}_{Y''}(-2mH')$  is torsion free and weakly positive. Thus there exists an integer  $n \in \mathbb{Z}_{>0}$  such that

$$\hat{\mathcal{S}}^n g'_*(\omega) \otimes \mathcal{O}_{Y'}(-2mnH') \otimes \mathcal{O}_{Y'}(nmH')$$

is generically generated by global sections for some  $n > 0$ , where  $\hat{\mathcal{S}}^n$  denotes the reflexive hull of  $\mathcal{S}^n$ . This implies that there is a divisor  $B'$  on  $Z'$  with  $\text{codim}'(B') \geq 2$  and an inclusion

$$\mathcal{O}_{Z''} \rightarrow \omega^n \otimes g'^*(-mnH') + B'$$

Therefore

$$M = mn(\sum n_{k'} E'_{k'} - g'^* p^*(K_X) - g'^*(H') + q^*(\delta L) - V + G) + B'$$

is effective (see [19]).

On the other hand, we can choose a family of general complete intersection curves  $C$  on  $Z'$  such that  $C$  does not intersect with the exceptional locus of  $Z' \rightarrow X$  (such as  $E'_{k'}, G$  and  $B'$ ), e.g., pull-back the general complete intersection curves from  $X$ .

Thus we have  $C \cdot g'^* p^*(K_Y) \leq C \cdot (q^*(\delta L) - V)$ . i.e., there exists a covering family of curves  $C'$  on  $Z$  such that  $C' \cdot g^* K_Y \leq C' \cdot (\delta L - q_* V)$ .

Let  $\delta \rightarrow 0$ , we get  $C' \cdot g^* K_Y \leq 0$ . If  $C' \cdot g^* K_Y < 0$ ,  $Y$  is uniruled [13]. Let us assume that  $C' \cdot g^* K_Y = 0$  (in particular,  $(\pi \circ q)_* V = 0$  and  $f$  is semistable in codim 1), and  $Y$  is not uniruled.

By a result in [2, Corollary 0.3],  $K_Y$  is pseudo-effective. Let  $K_Y \equiv P + N$  be the divisorial Zariski decomposition of  $K_Y$ , where  $P$  is the positive part (nef in codimension 1) and  $N$  is the negative part (an effective  $R$ -divisor) [1], [14]. Since the family of the curves  $g(C')$  also forms a *connecting* family [2] and  $g(C') \cdot K_Y = 0$ , we have  $p \equiv 0$  [2, Theorem 9.8]. Thus  $\kappa_\sigma(Y) = 0$  and hence  $\kappa(Y) = 0$  by a result of N. Nakayama [14, Proposition 6.2.8]. q.e.d.

*Proof of Corollary 1.* Let  $C$  be a general complete intersection curve on  $X$ . If  $C \cdot (K_X + D) < 0$ , then  $X$  is uniruled and there exists a non-trivial maximal rationally connected fibration  $f : X \dashrightarrow Y$  [3], [11]. By a result in [7],  $Y$  is not uniruled if  $\dim Y > 0$ . If  $Y$  is a point, then  $X$  is rationally connected. If  $\dim Y > 0$ , then  $\kappa(Y) = 0$  by the Main Theorem.

So let us assume that  $C \cdot (K_X + D) = 0$  and  $X$  is not uniruled. Thus we have  $D = 0$  and  $K_X$  is numerically trivial. Let  $\pi : Z \rightarrow X$  be a log resolution. Then  $K_Z + \Delta = \pi^* K_X + E$ , where  $\Delta$  and  $E$  are effective and  $\pi$ -exceptional. Let  $C'$  be the family of the curves from the pull-back of  $C$ . Then

$C' \cdot (K_Z + \Delta) = 0$ . Since  $K_Z$  is pseudo-effective, we have  $K_Z = P + N$  (the divisorial Zariski decomposition of  $K_Z$ ). Since  $\text{Supp}(\Delta)$  is  $\pi$ -exceptional,  $P$  is also the positive part of the divisorial Zariski decomposition of  $K_Z + \Delta$ . On the other hand,  $C$  forms a connecting family. Therefore we have  $P \equiv 0$  and hence  $K_\sigma(Z) = \kappa(Z) = 0$  [14, Proposition 6.2.8]. q.e.d.

*Proof of Corollary 2.* Since for any subvariety  $W$  of an abelian variety,  $\kappa(W) \geq 0$  and  $\kappa(W) = 0$  if and only if  $W$  is an abelian variety. q.e.d.

*Proof of Corollary 3.* We keep the same notations as in the proof of the Main Theorem. Let  $\pi : Z \rightarrow X$  be a log resolution of  $X$  and  $K_Z = \pi^*(K_X + D) + \sum e_i E_i$ , where  $e_i \geq -1$ . We may assume that  $g = f \circ \pi$  is a morphism.

As before, we have

$$K_{Z/Y} + \sum \epsilon_i E_i + \delta L \sim_Q \pi^*(K_X + D) + \sum m_i E_i - g^* K_Y + \delta L$$

where  $L$  is an ample divisor on  $Z$ .  $\epsilon_i = -e_i$  if  $e_i < 0$  and  $\epsilon_i = \{-e_i\}$  if  $e_i \geq 0$ , and  $m_i = \lceil e_i \rceil$  (for  $e_i \geq 0$ ). Since the nef reduction  $f$  is an almost holomorphic map [5], we have  $\pi^*(K_X + D)|_F \equiv 0$  (" $\equiv$ " means *numerically equivalent*) for the general fibers  $F$  of  $g$ . Thus

$$g_*(m(\pi^*(K_X + D) + \sum m_i E_i - g^* K_Y + \delta L)) \neq 0$$

for  $m \gg 0$ . By Lemma 1, we may assume that

$$m(\pi^*(K_X + D) + \sum m_i E_i - g^* K_Y + \delta L) + B$$

is an effective divisor for  $m \gg 0$ , where  $B$  is  $g$ -exceptional. Again as before, we can choose a covering family of curves  $C$  on  $Z$ . Since  $\dim g(C) > 0$ , we have  $C \cdot \pi^*(K_X + D) < 0$  by the general properties of nef reductions [5]. Let  $\delta \rightarrow 0$ , we have  $g(C) \cdot K_Y < 0$  and hence  $Y$  is uniruled. q.e.d.

**Acknowledgment:** I would like to thank S. Boucksom and F. Campana for helpful communications. I am grateful to J. Kollár for stimulating discussions and encouragement. I am also grateful to Y. Kawamata and E. Viehweg for the valuable advice and suggestions.

## REFERENCES

- [1]. S. Boucksom, *On the volume of a big line bundle*, Intern. J. Math **13** (2002), 1043-1063.
- [2]. S. Boucksom, J. Demailly, M. Paun, T. Peternell, *The pseudo-effective cone of a compact Kähler manifolds and varieties of negative Kodaira dimension*, math.AG/0405285 (2004).

- [3]. F. Campana, *Connexité rationnelle des variétés de Fano*, Ann. Sci. E.N.S **25** (1992), 539-545.
- [4]. F. Campana, *Orbifolds, Special Varieties and Classification Theory*, Ann. Inst. Fourier, Grenoble **54**(3) (2004), 499-630/math.AG/0110051(2001).
- [5]. T. Bauer, F. Campana, T. Eckl, S. Kebekus, T. Peternell, S. Rams, T. Szemberg, L. Wotzlaw, *A reduction map for nef line bundles*, Collection of papers dedicated to Hans Grauert, Springer-Verlag (2002), 27-36.(math.AG/0106147).
- [6]. J. Demailly, T. Peternell, M. Schneider, *Compact Kähler manifolds with Hermitian semi-positive anticanonical bundle*, Compositio. Math **101** (1996), 217-224.
- [7]. H. T. Graber, J. Harris, J. Starr, *Families of rationally connected varieties*, J. Amer. Math. Soc. **16** (1) (2003), 57-67.
- [8]. Y. Kawamata, *Characterization of abelian varieties*, Comp. Math. **43** (1981), 253-276.
- [9]. Y. Kawamata, *Subadjunction of log canonical divisors. II*, Amer.J. Math. **120** (5) (1998), 893-899.
- [10]. Y. Kawamata, K. Matsuda, K. Matsuki, *Introduction to the minimal Model Problem*, Adv. Stud. Pure. Math. **10** (1987), 283-360.
- [11]. J. Kollár, Y. Miyaoka, S. Mori, *Rationally connected varieties*, J. Alg. Geom. **1** (1992), 429-448.
- [12]. S. Lu, *A refined Kodaira dimension and its canonical fibration*, math.AG/0211029 (2002).
- [13]. Y. Miyaoka, S. Mori, *A numerical criterion for uniruledness*, Ann. of Math. **124** (1986), 65-69.
- [14]. N. Nakayama, *Zariski-Decomposition and Abundance*, RIMS-1142 (1997).
- [15]. T. Peternell, F. Serrano, *Threefolds with numerically effective anticanonical class*, Collectanea Math **49** (1998), 465-517.
- [16]. H. Tsuji, *Numerical trivial fibration*, math.AG/0001023 (2000).
- [17]. E. Viehweg, *Weak positivity and the additivity of Kodaira dimension for certain fiber spaces*, Adv. Stud. Pure. Math **1** (1983), 329-353.
- [18]. Q. Zhang, *On projective manifolds with nef anticanonical bundles*, J. reine angew. Math. **478** (1996), 57-60.
- [19]. Q. Zhang, *Rational connectedness of log  $Q$ -Fano varieties*, math.AG/0408301 (to appear in J. reine angew. Math).
- [20]. J. McKernan, Y. Prokhorov, *Threefold Thresholds*, Manuscripta. Math **114** (2004), 281-304/math.AG/0304152(2003).
- [21]. T. Bauer, T. Peternell, *Nef reduction and anticanonical bundles*, math.AG/0310484 (2003).

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA, MO 65211.  
U.S.A

*E-mail address:* `qi@math.missouri.edu`